

The Impact of Sleep Quality and Heart Rate Variability on Suicidal Ideation and Behavior in Depressed Patients: A Review

Alimire Alimu^{1,2*}, Yi Zhang²

¹ Graduate School, Xinjiang Medical University, Urumqi, China

² Department of Clinical Psychology, Xinjiang Uygur Autonomous Region People's Hospital, Urumqi, China

Email Address

515337975@qq.com (Yi Zhang)

*Correspondence: 515337975@qq.com

Received: 16 March 2024; **Accepted:** 12 April 2024; **Published:** 16 May 2024

Abstract:

This review sought to examine the relationship between sleep quality, heart rate variability (HRV), suicidal ideation, and behavior in patients with depression. An in-depth review of current literature revealed a consistent correlation between disrupted sleep patterns, altered heart rate variability, and increased suicidal tendencies in depressed patients. Compromised sleep quality, evident in insomnia and frequent nighttime awakenings, is a common symptomatic feature of depression and has been identified as a potential predictive factor for suicidality. Moreover, decreased HRV reflects impaired autonomic regulation, often linked with emotional dysregulation and increased suicide risk. Biopsychosocial factors and pathways that could potentially connect sleep disturbances and HRV to suicidal ideation and behavior are discussed, along with an emphasis on the importance of early detection and intervention in the potential mitigation of suicide risk in this vulnerable population. Further research is necessary to better understand these associations and translate findings into effective clinical interventions. Through increased comprehension of the interplay between these variables, the treatment strategies for depressed patients may be enhanced and the risk of suicidal ideation and behavior mitigated.

Keywords:

Depression, Sleep, Heart Rate Variability, Suicidal Ideation, Suicidal Behavior

1. Introduction

The significance of quality sleep and regulated HRV in maintaining optimal mental health is undeniably salient. Individuals experiencing Depressed Patients often grapple with altered sleep patterns, frequent awakenings at night, and perturbation in their heart rate variability, factors that contribute to increased susceptibility towards suicidal thoughts and behavior.

Depression, a debilitating mental disorder characterized by pervasive melancholy, hopelessness, fatigue, and a myriad of other symptoms, is recognized globally as a

significant public health issue [1]. It has been estimated to affect over 300 million people worldwide, leading to substantial impairment and disability. Suicidal ideation and behavior, two of the most severe outcomes of depression, represent a significant global health concern, contributing to the mortality rate associated with the disorder [2]. The relationship between the sleep quality, heart rate variability, and suicidal ideation and behavior in patients with depression is a complex but crucial field of study requiring exploration. The impact of diminished sleep quality alongside impaired heart rate variability on suicidal ideation and behavior cannot be underestimated in depression management. Each concept – Sleep Quality, Heart Rate Variability, Suicidal Ideation, Behavior, and Depression presents a unique influence on patients' mental health and can act as potential indicators or predictors for suicidality.

This review aspires to shed light on the intricate relationships and interactions between these key concepts within the field of mental health. Not only does this warrant a thorough synthesis of the current literature to delineate clear associations but also underlines the urgency to highlight the role of early detection and interventions. The deliverables from this review aim to contribute substantially to enhance clinical practice by informing better treatment strategies for depressing patients while mitigating the risk of suicidal ideation and behavior.

1.1. Definition of key concepts: Sleep Quality, Heart Rate Variability, Suicidal Ideation, Behavior, and Depression

Sleep Quality is an integral aspect of an individual's mental and physical health that can be categorized into three components: duration, continuity, and depth. It is compromised in cases of sleep disorders like insomnia characterized by difficulty falling asleep, staying asleep, or frequent nighttime awakenings [3].

HRV, a physiological phenomenon, represents the time variation between consecutive heartbeats. It serves as a biomarker of the body's adaptive response to stress, with decreased HRV often signaling a dysregulated function of the autonomic nervous system. HRV is greatly influenced by the Autonomic Nervous System(ANS), and lower HRV levels are associated with compromised ANS function, which can manifest in emotional dysregulation and other symptoms [4].

Suicidal Ideation refers to thinking about, considering, or planning suicide. It is a critical warning sign in mental health disorders, specifically depression, marking a heightened level of distress and the need for immediate intervention [5].

Behavior, in this context, refers to the observable conduct of an individual, including their interactions with self, others and environmental stimuli. In the realm of mental health, behavior includes overt manifestations of mental distress such as self-harm or suicidal attempts.

Depression, officially termed Major Depressive Disorder (MDD), is a common and debilitating mood disorder characterized by persistent feelings of sadness, loss of interest in activities previously enjoyed, difficulty concentrating, and changes in sleep patterns among other symptoms. It is often associated with suicidal ideation and behavior.

Understanding the interlink between these concepts is essential for elucidating the complex cognitive, physiological, and social mechanisms underlying depression and suicide, and hence this review.

1.2. An overview of depression and suicidal ideation and behavior

Studies have shown that a majority of patients with depression report a diminishment in both sleep quality and quantity, primarily characterized by insomnia and frequent awakenings during the night. This incurs a detrimental impact on daytime functioning, leading to increased feelings of fatigue and despondency [6]. Moreover, disturbed sleep patterns have been consistently linked with an increased risk of suicidal ideation and behavior in depressed patients.

Suicidal ideation refers to thoughts about self-harm or a wish to die, with or without a specific suicide plan. Suicidal behavior, on the other hand, encompasses a spectrum of actions that range from non-fatal self-harm to completed suicide. A substantial body of research has demonstrated a strong interconnectedness between depression, suicidal ideation, and suicide attempts. This disconcerting association is underscored by the fact that suicide is now one of the leading causes of death among individuals with MDD.

The role of HRV in this intricate relationship between depression and suicidality is of growing interest in the field of mental health. Accumulating evidence points towards a link between attenuated HRV and an elevated risk for suicide amongst depressed patients [7].

In sum, this brief overview underscores the importance of exploring the multi-faced relationship between sleep disturbances, altered HRV, and suicidal tendencies in patients with depression. Understanding this connection has the potential to illuminate crucial predictive factors for suicidality and guide more targeted interventions to mitigate suicide risk in this vulnerable population.

1.3. The purpose of the review and its relevance in the field of mental health

This review aims to analyze and synthesize evidence linking sleep quality and HRV to suicidal ideation and behavior in patients with depression. The goal is to deepen understanding of these associations and highlight potential preventive and intervention strategies to reduce suicide risk in this patient population. Suicidal ideation and behavior in depressed patients pose a significant challenge in mental health, and early detection and intervention based on factors like sleep quality and HRV could be critical in saving lives. Exploring the relationships between these variables could help mental health professionals identify high-risk individuals more accurately and tailor interventions effectively to prevent suicide. In this light, this review is highly pertinent to the field of mental health, as it seeks to provide a conceptual framework that could guide research, clinical practice, and policy-making towards improved suicide prevention in patients with depression. Future studies should prioritize investigating the underlying biopsychosocial mechanisms connecting sleep quality and HRV to suicidal ideation and behavior, enriching our knowledge base and empowering mental health specialists with evidence-based practices to address this important public health issue.

2. Sleep Quality and Suicide

Sleep quality plays a significant role in maintaining optimal mental health, where disturbances to regular sleep patterns often predicate the onset of psychiatric disorders. Normal sleep architecture purportedly aids in emotional regulation and the proliferation of neurons within the brain. Conversely, deprived or fragmented sleep can lead to impairments in cognitive functioning, emotional stability, and overall

mental well-being [8]. Specifically concerning depression, poor sleep quality is not merely a symptomatic marker but also a contributing factor. Quantitative literature investigating the sleep-depression dyad has established a bidirectional relationship between the two [9]. Insomnia and frequent awakenings during the night, symptomatic of poor sleep quality, are common among depressed patients. Insomniac patients carry a tenfold risk of developing depression compared to those without sleep disturbances. The altered circadian rhythm perpetuates the depressive state by interfering with normal neurochemical processes, further compromising sleep quality and instigating a vicious cycle [10,11]. Sleep problems are more common among depressed patients with suicidal ideation compared to non-suicidal depressed patients. Specifically, insomnia, nightmares, and decreased sleep efficiency are associated with an accentuated risk of suicidal ideation and attempts. Moreover, sleep disturbances contribute to hopelessness - a robust predictor of suicide - reinforcing the linkage between poor sleep quality and suicidal tendencies.

There is accumulated evidence to suggest that poor sleep quality may serve as a risk factor for suicidal ideations and behaviors, and not merely a symptom of a depressive state. However, further comprehensive research is needed to validate these findings and to explore the precise underlying mechanisms that link poor sleep quality, depression, and suicide.

2.1. The role of sleep quality in mental health

Sleep quality is an integral part of overall mental health. It has been widely studied and firmly established that adequate, restful sleep is fundamental to overall well-being and directly impacts cognitive function, emotional regulation, and physical health. This section sheds light on how disrupted sleep patterns adversely affect mental health, amplifying the existing psychological distress in individuals.

Lack of sleep or poor quality of sleep has been associated with various mental health disorders. It can decrease an individual's resistance to stressful situations, heighten the emotional responses, and weaken cognitive function which interacts negatively with both the psychological and biological aspects of mental health. Insomnia, a primary indicator of poor sleep quality, has been linked to the onset and exacerbation of several psychiatric conditions such as anxiety disorders, depression, and bipolar disorder [12].

More specific to the context of the present review, disrupted sleep patterns, seen in the form of compromised sleep quality, are a common symptomatic feature of depression. It is worth noting that these sleep disruptions not only function as an indicator or symptom of a pre-existing condition but have also been shown to play a significant role in the development, recurrence, and severity of mental health conditions, including depression.

Research shows that the underlying mechanisms of sleep's effect on mental health are multifaceted. On a biological level, lack of sleep causes imbalances in various neurotransmitters and stress hormones, thus aggravating mood disorders [13,14]. Sleep disruption is similarly believed to diminish the efficacy of the pre-frontal cortex, an area of the brain responsible for tasks related to emotional regulation.

In summary, sleep quality substantively intersects with mental health, with poor sleep quality contributing to the onset and severity of mental health disorders. Its impact, as reviewed above, is increasingly noticeable in conditions like depression, warranting further exploration in the context of suicidal ideation and behavior within

this review. The following sections will delve further into the relationship between poor sleep quality, depression, and suicidal ideation and behavior.

2.2. The relationship between poor sleep quality and depression

A substantial body of research has identified a consistent and remarkable relationship between poor sleep quality and depression. This relationship is characterized by a multidimensional interaction, underpinning depression as both a consequence and a contributor to poor sleep quality.

Depressive disorder often co-exists with significant sleep disturbances [15]. These disturbances can manifest in various ways including difficulties in falling asleep, repeated nocturnal awakenings, and early-morning awakenings, or even hypersomnia. Epidemiological studies consistently suggest that sleep problems, particularly insomnia, are prevalent in patients grappling with depression. As high as 90% of people suffering from depression report experiencing troubles with their sleep, illustrating the close connection between sleep quality and depression [16].

The relationship between these two entities is a reciprocal one. On the one hand, poor sleep quality can exacerbate depressive symptoms, creating a cycle of worsening mental health. People experiencing sleep disturbances tend to show a pessimistic outlook, impaired cognitive functioning, and decreased quality of life, all of which can lead to or aggravate depression. On the other hand, depression itself can lead to poor sleep quality, a fact that is reflected in disturbed sleep patterns becoming a criterion included in the diagnostic symptomatology for major depressive disorder.

Neurobiological factors and processes are also likely to play a significant role in the relationship between sleep quality and depression. Similar physiological pathways may be affected in both depression and sleep disturbances, such as alterations in circadian rhythms, dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis, and anomalies in brain neurotransmitters [17]. Therefore, alterations in these physiological systems manifesting as sleep disturbances may also be implicated in the pathogenesis of depression.

2.3. Impact of sleep disturbances on suicidal ideation and behavior in depressed patients

Sleep disturbances are a significant factor in suicidal ideation and behaviour amongst depressed patients. There appears to be a consistent correlation between disrupted sleep patterns and increased suicidal tendencies for those suffering from depression. The most common symptomatic evidence often revolves around compromised sleep quality, characterized by insomnia and frequent nighttime awakenings, and these have been recurrently flagged as potential predictive factors for suicidality [11].

Depressed patients with sleep disturbances have been observed to display exaggerated depressive symptoms, with escalated dimensions of hopelessness and cognitive distortions, often culminating in suicidal ideation which, unfortunately, could be a precursor for suicide attempts. The correlation between sleep disturbances and suicidal tendencies is multidimensional inherently- both directly and as a magnifying factor of depressive symptoms. On one hand, sleep disturbances potentially illuminate core processes exacerbating depression and suicidality including impaired cognitive functioning, and alteration of emotional processing and regulation that accentuates negative emotions [18].

On another outlook, sleep disturbances in depressed patients might indirectly enhance suicide risk by virtue of potentiating other well-established suicide risk factors such as psychosocial stressors, impulsivity, or alcohol and substance abuse which are frequently co-occur with disturbed sleep [19].

Furthermore, it is also worth noting that, reciprocal relationships likely exist, strengthening the sinister intersection of sleep disturbances, depression, and suicidality. For instance, suicidal ideation or behaviors could further disrupt the already vulnerable sleep of depressed individuals and erode their sleep quality, initiating a possible vicious cycle.

Moreover, the entirety of these suggestions bring to the forefront the necessity for proper early detection and intervention methodology, paramount in the potential mitigation of suicide risk in this vulnerable population. Through a comprehensive understanding of these interlinkages, treatment strategies for depressed patients could be enhanced and the risk of suicidal ideation and behavior may be mitigated.

Lastly, it is crucial to acknowledge the need for further research to fortify these theoretical claims in strengthening our understanding of this relationship, facilitating developments in relevant clinical interventions. Through increased comprehension of the interplay between these variables, the treatment strategies for depressed patients may be enhanced and the risk of suicidal ideation and behavior mitigated.

3. HRV and Suicide

Research suggests that individuals with high HRV often exhibit strong emotional regulation and resilience to psychological stress [20]. Conversely, a decrease in HRV is seen as a biomarker for various psychological conditions, such as depression. Low HRV indicates a reduced capacity for autonomic regulation, which is closely linked to emotional dysregulation and can lead to higher levels of psychological distress. Impaired autonomic control, often seen in low HRV, is associated with an increased risk of severe mental health issues, including suicidal ideation and behavior. Studies have shown a strong connection between low HRV and suicidal tendencies in patients with depression, highlighting the importance of HRV in suicide risk assessment and prevention strategies [21]. Further research is needed to better understand these relationships and develop effective clinical interventions.

3.1. Understanding HRV and its significance in mental health

The complex and diverse biological mechanisms of suicidal behavior in depressed individuals often involve low HRV, which is considered a potential predictor of suicide risk [22]. Utilizing 24-hour ambulatory electrocardiogram monitoring allows for a comprehensive evaluation of HRV levels in patients, shedding light on heart rate fluctuations at rest and in response to stress. These insights collectively support the idea that a significant correlation exists between decreased HRV and suicide risk in depressed patients, particularly when the LF power is elevated, indicating a higher predictive value for suicide risk assessment [23]. Given these compelling correlations, healthcare professionals should recognize the importance of HRV as an indicator of psychological stress in preventing suicidal behaviors. It is essential to tailor individualized psychological interventions and management strategies for patients by integrating HRV data with other clinical information, such as insomnia status, pain experience, and depression severity. By combining physiological indicators with clinical data and employing scientific statistical models, a quantitative foundation for

assessing suicide risk in depressed patients can be established, facilitating early identification and intervention.

3.2. Connection between low HRV and emotional dysregulation

models of mental health, HRV emerges as a crucial physiological marker for emotional regulation. A substantial body of research has established a connection between decreased HRV and emotional dysregulation, a key characteristic often observed in individuals with depressive disorders. Emotional dysregulation, marked by intense and poorly managed emotional reactions, can impede a person's ability to handle life's challenges in a balanced and adaptive manner [24]. Essentially, HRV reflects the autonomic nervous system's health and its ability to adjust to environmental stimuli, which is closely linked to emotional regulation [26]. Lower HRV indicates a less adaptive autonomic nervous system, typically associated with increased emotional inflexibility and less effective emotion-regulation strategies. Numerous scientific studies have supported this concept, with research showing reduced HRV in patients diagnosed with various affective disorders, indicating compromised autonomic function and reduced emotional adaptability. This manifests as maladaptive emotional responses, persistent feelings of hopelessness, irritability, and struggles with impulse control [25]. Moreover, the correlation between low HRV and emotional dysregulation has direct implications for suicide risk among individuals with depression. Given that emotional dysregulation is linked to ineffective coping mechanisms, it contributes to suicidal thoughts and behaviors. Decreased HRV, reflecting impaired emotion regulation, signifies a reduced ability to cope with emotional distress, thereby increasing the likelihood of suicidal tendencies in depressed patients.

3.3. The correlation between altered HRV and suicide risk in depressed patients

HRV, as previously discussed, is an index of the body's autonomic regulation capability which, when impaired, can contribute to emotional dysregulation with evidence suggesting a direct connection to heightened suicide risk.

Studies on patients diagnosed with Major Depressive Disorder have shown a noticeable decrease in HRV, indicating compromised parasympathetic function and subsequent dysregulation in emotional responses. This dysregulation is believed to play a significant role in the development of suicidal behavior. Additionally, concurrent sleep disturbances in individuals with depression, such as insomnia and frequent nighttime awakenings, may worsen this effect, as sleep plays a restorative role in autonomic and emotional regulation systems. Some researchers suggest that HRV could be a reliable biological marker for assessing suicide risk. Changes in HRV are strongly linked to suicidal thoughts and actions, with depressed patients who have attempted suicide often exhibiting lower HRV compared to those who have not [28]. Furthermore, a decline in HRV frequently precedes suicide attempts, indicating its potential predictive value. It is important to note, however, that while substantial evidence supports the connection between low HRV and increased suicidality, the exact nature of this relationship remains somewhat unclear, highlighting the need for further investigation [27]. The prospect of low HRV not only being a consequence but also a precursor of suicide risk emphasizes the importance of early HRV assessment in clinical settings for individuals with depression. This approach shows promise for early detection and intervention, contributing to the overarching goal of preventing self-harm and suicide in this vulnerable population.

4. Interplay between Sleep Quality, Heart Rate Variability, and Suicide

The intricate connection between sleep quality, HRV and suicidality in depressed patients forms a complex framework which requires a meticulous examination. Sleep disturbances, involving insomnia and frequent awakenings during the night, are commonly observed in the depressed population and have been linked to increased suicidal ideation and behavior [29]. Altered HRV has also emerged as an important physiological consequence in depression, reflecting a dysfunctional autonomic regulation, which in turn intensifies the risk of suicide into a concrete reality.

At the crux of this trifaceted interplay, there is an array of biopsychosocial influences. Biological factors are prevalent, with sleep deprivation impairing neurocognitive functions, heightening emotional response, and contributing to disrupted mood regulation, known precursors to suicidal ideation [30]. Similarly, decreased HRV is often coupled with other physiological abnormalities like increased inflammatory response and HPA axis dysregulation, known to exacerbate depressive symptoms.

Social factors come into play as worsening sleep quality and increased social isolation often display a reciprocal relationship. The deterioration of sleep quality can induce feelings of loneliness, leading to social withdrawal, ultimately amplifying suicidal ideation [31]. Psychological factors also play a significant role, with chronically disturbed sleep patterns and altered HRV often leading to heightened feelings of hopelessness and helplessness, key markers for suicidal behavior.

In conclusion, the entwined relationship of sleep disturbances, altered HRV, and suicidal ideation in the depressed population warrants focused attention. A comprehensive understanding of this triad could potentially enhance treatment strategies, justify early detection strategies, and ultimately mitigate the risk of suicide in this vulnerable group [32]. However, further research is still needed to fully unveil the deep-seated interlinks.

5. Conclusive Remarks and Future Directions

The current review unveiled a consistent correlation between sleep disorders, altered HRV, and an increased proclivity towards suicidal ideation among patients with depression. These established connections suggest that these variables both independently contribute and synergistically interact to shape the suicide risk landscape among mentally afflicted patients.

Compromised sleep quality, particularized by phenomena such as insomnia and frequent nocturnal arousals, has been recurrently recognized as a symptomatic hallmark of depressive disorders. Emerging research narratives have suggested that these sleep disturbances might not merely be an accompanying feature but might play a pivotal role in precipitating suicidal tendencies, emphasizing the need for a more in-depth investigation into delineating exact mechanisms that underpin this association.

Similarly, the compilation of current literature also highlighted that depressed patients often exhibited diminished HRV, indicative of a disrupted autonomic regulatory system. This dysregulation is frequently correlated with heightened emotional instability, further accentuating suicide susceptibility. Therefore, HRV could offer a promising marker for diagnosis and risk assessment.

Early detection and timely intervention could significantly revolutionize the treatment outcomes for depressive patients. Mapping sleep patterns and tracking HRV could enable healthcare providers to proactively identify patients at increased suicide risk, allowing for necessary treatment modifications or interventions.

Future explorations should commit to constructing a robust evidence base through large-scale, longitudinal studies with more diverse samples to validate these associations. The impact of potential confounding factors such as socioeconomic influences, lifestyle changes, and comorbid psychiatric conditions should also be considered.

Ultimately, these findings could potentially transform clinical practices by offering an evidence-based intervention strategy. A more comprehensive understanding of the role of sleep and HRV in suicide risk can help tailor personalized treatment plans for depressed patients, addressing these anomalies and thereby mitigating the likelihood of suicide.

In conclusion, understanding the complex interplay of sleep quality, HRV, and suicidality can move us one step closer in not only improving the management of depression but also in working towards the prevention of suicide in a high-risk population.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this article.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

References

- [1] American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-51). USA: American Psychiatric Publication; 2013 (n.d.).
- [2] Mirchandaney, R.; Asarnow, L.D.; Kaplan, K.A. Recent advances in sleep and depression. *Curr Opin Psychiatry*, 2023, 36, 34-40.
- [3] Riemann, D.; Krone, L.B.; Wulff, K.; Nissen, C. Sleep, insomnia, and depression. *Neuropsychopharmacology*, 2020, 45, 74-89.
- [4] Provini, F. Sleep disorders and the autonomic nervous system. *J Neurol Sci*, 2021, 429: 117905.
- [5] Zhao, H.; Jiang, C.; Zhao, M.; Ye, Y.; Yu, L.; Li, Y., et al. Comparisons of Accelerated Continuous and Intermittent Theta Burst Stimulation for Treatment-Resistant Depression and Suicidal Ideation. *Biol Psychiatry*, 2023, S0006322323017882.
- [6] Bodicherla, K.; Tubbs, A.; Fernandez, F.X.; Perlis, M.; Grandner, M. 776 Perceived sleep control and nightmares distinguish college students with suicidal ideation from past attempters. *Sleep*, 2021, 44, A302-A302.
- [7] Rüesch, A.; Villar, De Araujo, T.; Bankwitz, A.; Hörmann, C.; Adank, A.; Ip, C.T. et al. A recent suicide attempt and the heartbeat: Electrophysiological

- findings from a trans-diagnostic cohort of patients and healthy controls. *J Psychiatr Res*, 2023, 157, 257-263.
- [8] Chen, R.F.; Cai, Y.; Zhu, Z.H.; Hou, W.L.; Chen, P.; Wang, J, et al. Sleep disorder as a clinical risk factor of major depression: associated with cognitive impairment. *Asian J Psychiatry*, 2022, 76, 103228.
- [9] Liu, H. Association between sleep duration and depression: A Mendelian randomization analysis. *J Affect Disord*, 2023, 335, 152-154.
- [10] Hassamal, S. Chronic stress, neuroinflammation, and depression: an overview of pathophysiological mechanisms and emerging anti-inflammatories. *Front Psychiatry*, 2023, 14, 1130989.
- [11] Chen, S.J.; Zhang, J.H.; Li, S.X.; Tsang, C.C.; Chan, K.C.C.; Au, C.T.; et al. The trajectories and associations of eveningness and insomnia with daytime sleepiness, depression and suicidal ideation in adolescents: A 3-year longitudinal study. *J Affect Disord*, 2021, 294, 533-542.
- [12] Diaz, A.P.; Fernandes, B.S.; Quevedo, J.; Sanches, M.; Soares, J.C. Treatment-resistant bipolar depression: concepts and challenges for novel interventions. *Braz J Psychiatry*, 2022, 44, 178-186.
- [13] Hu, X.; Cheng, L.Y.; Chiu, M.H.; Paller, K.A. Promoting memory consolidation during sleep: A meta-analysis of targeted memory reactivation. *Psychol Bull*, 2020, 146, 218-244.
- [14] Hopkins, S.C.; Dedic, N.; Koblan, K.S. Effect of TAAR1/5-HT1A agonist SEP-363856 on REM sleep in humans. *Transl Psychiatry*, 2021, 11, 228.
- [15] Hein, M.; Lanquart, J.P.; Loas, G.; Hubain, P.; Linkowski, P. Prevalence and risk factors of excessive daytime sleepiness in major depression: A study with 703 individuals referred for polysomnography. *J Affect Disord*, 2019, 243, 23-32.
- [16] Riemann, D.; Benz, F.; Dressle, R.J.; Espie, C.A.; Johann, A.F.; Blanken, T.F. et al. Insomnia disorder: State of the science and challenges for the future. *J Sleep Res*, 2022, 31, e13604.
- [17] Drinčić, T.; Van Dalen, J.H.; Kamphuis J.; Jentsch, M.C.; Van Belkum, S.M.; Meddens, M.J.M, et al. The Relationship between Insomnia and the Pathophysiology of Major Depressive Disorder: An Evaluation of a Broad Selection of Serum and Urine Biomarkers. *Int J Mol Sci*, 2023, 24, 8437.
- [18] Caudle, M.M.; Dugas, N.N.; Patel, K.; Moore, R.C.; Thomas, M.L.; Bomyea, J. Repetitive negative thinking as a unique transdiagnostic risk factor for suicidal ideation. *Psychiatry Res*, 2024, 334, 115787.
- [19] Pitman, A.; McDonald, K.; Logeswaran, Y.; Lewis, G.; Cerel, J.; Lewis, G.; Erlangsen, A. The role of depression and use of alcohol and other drugs after partner suicide in the association between suicide bereavement and suicide: cohort study in the Danish population. *Psychol Med*, 2024, 1-10.
- [20] Čukić, M.; Savić, D.; Sidorova, J. When Heart Beats Differently in Depression: Review of Nonlinear Heart Rate Variability Measures. *JMIR Ment Health*, 2023, 10, e40342.
- [21] Wilson, S.T.; Chesin, M.; Fertuck, E.; Keilp, J.; Brodsky, B.; Mann, J.J. et al. Heart rate variability and suicidal behavior. *Psychiatry Res*, 2016, 240, 241-247.

- [22] Department of Neuropsychiatry of Slovak Medical University in Bratislava and Psychiatric Hospital of Philippe Pinel in Pezinok, Pezinok, Slovakia, Rakus T, Third faculty of medicine Charles University in Prague, Czech republic, Hubcikova K, Department of Neuropsychiatry of Slovak Medical University in Bratislava and Psychiatric Hospital of Philippe Pinel in Pezinok, Pezinok, Slovakia, Department of Neuropsychiatry of Slovak Medical University in Bratislava and Psychiatric Hospital of Philippe Pinel in Pezinok, Pezinok, Slovakia, et al. *ELECTROPHYSIOLOGICAL CORRELATES OF SUICIDALITY. Psychiatr Danub*, 2021, 33, 266-279.
- [23] Lee, D.; Baek, J.H.; Cho, Y.J.; Hong, K.S. Association of Resting Heart Rate and Heart Rate Variability With Proximal Suicidal Risk in Patients With Diverse Psychiatric Diagnoses. *Front Psychiatry*, 2021, 12, 652340.
- [24] Chai, H.W.; Jester, D.J.; Lee, S.; Joo, S. LOSS, SLEEP, AND HEART RATE VARIABILITY: GENDER DIFFERENCES. *Innov Aging*, 2019, 3, S915-S916.
- [25] Gullett, N.; Zajkowska, Z.; Walsh, A.; Harper, R.; Mondelli, V. Heart rate variability (HRV) as a way to understand associations between the autonomic nervous system (ANS) and affective states: A critical review of the literature. *Int J Psychophysiol*, 2023, 192, 35-42.
- [26] Santo, F.D.; Velasco, Á.; Rodríguez-Revuelta, J.; González-Blanco, L.; Álvarez-Vázquez, C.M.; García-Portilla, M.P, et al. P.341 Childhood maltreatment in adult depressed patients: impact of types and number of traumas on suicidal behaviour. *Eur Neuropsychopharmacol*, 2019, 29: S243.
- [27] Adolph, D.; Teismann, T.; Forkmann, T.; Wannemüller, A.; Margraf, J. High frequency heart rate variability: Evidence for a transdiagnostic association with suicide ideation. *Biol Psychol*, 2018, 138, 165-171.
- [28] Kim, J.M.; Kim, J.W. Impact of suicidal ideation on long-term cardiac outcomes in patients with acute coronary syndrome. *IBRO Rep*, 2019, 6, S71.
- [29] Tubbs, A.S.; Khader, W.S.; Hale, L.; Branäs, C.; Perlis, M.; Gehrels, J.A. et al. 0926 Sleep Timing And The Prevalence Of Suicidal Ideation In A Community Sample. *Sleep*, 2019, 42: A372-A373.
- [30] Chen, H.; Hong, L.; Tong, S.; Li, M.; Sun, S.; Xu, Y, et al. Cognitive impairment and factors influencing depression in adolescents with suicidal and self-injury behaviors: a cross-sectional study. *BMC Psychiatry*, 2023, 23, 247.
- [31] Dorrian, J.; Centofanti, S.; Smith, A.; McDermott, K.D. Self-regulation and social behavior during sleep deprivation. *Progress in Brain Research*, 2019, 246. Elsevier, 73-110.
- [32] Gallo, J.J.; Hwang, S.; Truong, C.; Reynolds, C.F.; Spira, A.P. Role of persistent and worsening sleep disturbance in depression remission and suicidal ideation among older primary care patients: the PROSPECT study. *Sleep*, 2020, 43, zsaa063.



© 2024 by the author(s); licensee International Technology and Science Publications (ITS), this work for open access publication is under the Creative Commons Attribution International License (CC BY 4.0). (<http://creativecommons.org/licenses/by/4.0/>)